

Chen and Itoi Reply

Wei Chen

Department of Physics, University of North Carolina, Chapel Hill, NC 27599-3255

Chigak Itoi

*Department of Physics and Atomic Energy Research Institute, College of Science and Technology,
Nihon University, Kanda Surugadai, Chiyoda-ku, Tokyo 101, Japan*

PACS numbers: 11.15.Bt, 11.15.Pg

In a Comment [1] statements on our paper [2] are made with which we disagree.

It is argued in [1] that the model (3) can not be equivalent to (1) and (2), as the former has twice as many degrees of freedom as the latter. To clarify the issue, let us recall two key ingredients to claim the equivalence in [2] and a sequent work [3]: I) introducing the coherent states and integrating out the matter and CS fields, we obtain a continuum random walk expression for the partition function, *e.g.* eq. (9) in [2], in which the spin of matter field j appears as the coefficient of spin factor and the CS coefficient α as the coefficient of relative and self-energies; and II) there have been topological relations between the spin factor and relative and self-energies regularized by point splitting, eqs. (12) and (13). The expression and topological relations suggest that an integer (or odd-half-integer) part of α , n , can be absorbed by changing j . Then, as the spin j characterizes the degrees of freedom carried by the matter field, the latter may be changed when the spin- j matter is mapped to spin- $(j+n)$ matter and α is shifted to $(\alpha-n)$. This indicates that CS interactions affect the degrees of freedom of the matter field and whole system. After all, the equivalent CS models describe the same (anyon) system and have the same amount degrees of freedom. While it has been known in many (1+1) dimensional field theories that interactions change the degrees of freedom carried by fields, it seems for us to have seen a possibility for such an example in (2+1) dimensions.

It is pointless to require equivalent models to involve same divergences. However, as a matter of fact, the models (1), (2) and (3) all need use of ultraviolet regularization, including in calculating Π_o in (1), contrary to that argued in [1]. The author of [1] further picks up difference between one-loop corrections to $\Pi_o(p)$ calculated in a particular regularization in (1) and (2) to argue inequivalence. We have two remarks to make. First, it makes no sense to compare non-universal quantities such as $\Pi_o(0)$'s in different models. Since these are regularization dependent, they can take any values that symmetries allow. Though, it is tempted to check the local relevant operator content. Non-vanishing $\Pi_o(0)$'s and

$\Pi_e(0)$'s imply that the current-current correlations in the low energy limit induce a CS term and a Maxwell term in both (1) and (2). Secondly, the non-local terms from $\Pi_o(p \neq 0)$ and $\Pi_e(p \neq 0)$ are finite in both (1) and (2), as they should be. Nevertheless, one should be aware that for any given α , at least two of the three models are strongly coupled ones, and thus it is difficult to check non-perturbative results by means of perturbation, even qualitatively sometimes. In particular, it makes no judge to the equivalence to simply compare the values of those non-local terms obtained in perturbation at one loop.

To argue the models (1) and (2) dissimilar, the author of [1] takes $\alpha \rightarrow \infty$ and discusses some process, as if there existed no CS terms. It is inconsistent at all, as large α 's amount to strong CS couplings that can not be ignored. This is readily seen by rescaling $a_\mu \rightarrow \sqrt{4\pi\alpha}a_\mu$ for (1), for instance, and the CS fermion coupling has the strength $\sqrt{4\pi\alpha}$. Obviously, when $\alpha \rightarrow \infty$, perturbation around the free fermion theory is meaningless. An effective approach to handle strong CS interactions has been suggested in [2] and [3]. That is to trade the integer part of α for higher spin of matter fields, and then perturbative calculations in the remaining weak CS interaction could be good approximations. In this case, taking $\alpha \rightarrow \infty$, the models (1), (2) and (3) are now all describing a spin- ∞ field coupled to the external field C_μ , and are thus formally equivalent (though it is not clear to us whether the limit is well-defined). On the other hand, if taking weak coupling limit in one model, $\alpha \rightarrow 0$ for instance, the CS field is decoupled in (1), while CS interactions are considerably strong in (2) and (3).

In conclusion, the author of [1] seems to have overlooked the highly non-trivial CS couplings, and his hand-waving arguments are totally groundless.

The work of W.C. was supported in part by the U.S. DOE grant No. DE-FG05-85ER-40219.

-
- [1] C.R. Hagen, UR-1353/ER-40685-804, hep-th/9404130.
 - [2] W. Chen and C. Itoi, Phys. Rev. Lett. **72**, 2527 (1994).
 - [3] W. Chen and C. Itoi, IFT-488-UNC/NUP-A-94-3, hep-th/9404107.